



# Soyfoods for Infants, Children & Adolescents

2010 EDITION



Establishing healthful eating habits early in life is important for two reasons. First, childhood eating habits track into adulthood and changing adult dietary behavior is difficult.<sup>1-5</sup> Second, evidence suggests that healthy behaviors in childhood and adolescence can affect the risk of developing certain chronic diseases later in life.<sup>6-9</sup> For example, obesity in childhood is associated with an increased mortality from cardiovascular disease in adulthood, independent of adult weight.<sup>10</sup> Early lifestyle factors are also known to affect the likelihood of developing breast cancer during adulthood.<sup>11</sup> These observations are important given that 20 percent of U.S. children are overweight<sup>12</sup> and diseases once seen primarily in adults, such as hypertension<sup>3</sup> and Type 2 diabetes mellitus,<sup>14</sup> are increasingly common in childhood. It is also recognized that the beginning stages of chronic diseases, such as coronary heart disease, are already apparent in adolescents.<sup>15, 16</sup>

Given the importance of early-life dietary behavior, it is essential to understand how the nutritional attributes of soyfoods may impact the health of young people from infancy through the teenage years.

## Soy Infant Formula

Although breast milk is the ideal food for infants,<sup>17</sup> about one-third of women choose not to or cannot breastfeed. Of those who do, most switch to formula feeding at some point in the infant's first year.<sup>18</sup> Commercially-prepared, fortified infant formulas are appropriate to supplement or replace human milk during the first year of life. Cow's milk formula is the most commonly used product, but about 15 percent of infants are fed soy formula for some period of time.<sup>19</sup>

An allergy to milk protein is among the most common reasons for placing an infant on soy formula. There is clear evidence that soy

formula is hypoallergenic relative to cow's milk formulas.<sup>18</sup> However, because 10-14 percent of infants who are allergic to cow's milk formula are also allergic to soy formula, the American Academy of Pediatrics (AAP) suggests that many infants with documented cow's milk protein allergy should be switched directly to a hydrolyzed protein formula.<sup>19</sup> In contrast, an Australian panel of experts recently concluded that soy formula is an appropriate alternative for infants over six months old who demonstrated immediate food allergy to cow's milk and delayed reaction in the form of atopic eczema and other gastrointestinal syndromes.<sup>20</sup>

## Isoflavones in Diets of Infants Fed Soy Formula

An estimated 20 million people in the United States consumed soy formula during infancy since it first became commercially available in the 1960s.<sup>19</sup> Several cases of goiter were identified in the mid-1960s in infants using soy formula, but this problem was eliminated soon thereafter with the advent of iodine fortification of the formula.<sup>23-25</sup> Since then, no problems specifically related to soy formula use have been identified and research shows that infants fed soy formula grow and develop normally.<sup>19, 26-29</sup> All soy formulas are fortified with iodine, iron, methionine, carnitine and taurine and contain 20 percent more calcium and phosphorous than cow's milk formulas. However, soy formula may be contraindicated for infants with congenital hypothyroidism who require synthetic thyroid hormone.<sup>30</sup> This is because of evidence suggesting soy protein is one of a number of factors that may interfere with the absorption of thyroid medication.<sup>31</sup>

*All soy formulas are fortified with iodine, iron, methionine, carnitine and taurine, and contain 20 percent more calcium and phosphorous than cow's milk formulas.*

Despite its long history of use, soy formula has become controversial in recent years due to the naturally high isoflavone content of the soybean.<sup>32, 33</sup> Isoflavones, often referred to as phytoestrogens, exhibit estrogen-like effects under certain experimental conditions.<sup>34</sup> However, isoflavones are not the same as estrogen. Research with adults shows that many biological measures that are affected by the hormone estrogen are not affected by isoflavones.<sup>35-55</sup> Furthermore, soy protein, which is used in soy formula, is not the same as isoflavones. Most importantly, there is no clinical evidence in infants that soy formula consumption leads to adverse effects, and recent results of a unique study are quite reassuring in regard to any potential hormonal effects.<sup>26, 56, 57</sup>

In this study, breast buds, uterus, ovaries, prostate and testicular volumes were assessed by ultrasonography in 40 breast fed, 41 milk



*Soyfoods provide high-quality protein and are low in saturated fat.*

formula fed and 39 soy formula fed infants at four months of age.<sup>58</sup> In all cases soy formula fed infants were similar to breastfed or milk formula fed infants whereas unexpectedly, milk formula fed infants had greater mean ovarian volume and greater numbers of ovarian cysts per ovary than did breastfed infants. The clinical relevance of these findings is unclear. Long-term data are limited, but in one retrospective study, no meaningful differences in a host of biological parameters between adults fed soy formula or cow's milk formula as infants were noted.<sup>59</sup> Interestingly, results from a very small and very preliminary study found that girls fed soy formula as infants were 40-60 percent less likely to develop breast cancer as adults compared to women who were fed breast milk, cow's milk formula or a combination of both.<sup>60</sup>

*Clinical research shows that soy protein directly lowers cholesterol levels in children.*

A comprehensive review published in 2004 summarized views on the isoflavone content of soy formula with this statement: "The evidence from laboratories showing biological activities at doses or tissue concentrations relevant to soy-fed infants is difficult to reconcile with the long record of uneventful use of these formulas."<sup>61</sup> This sentiment is similar to the current position of the AAP, which was issued in 2008: "... although studied by numerous investigators in various species, there is no conclusive evidence from animal, adult human, or infant populations that dietary soy isoflavones may adversely affect human development, reproduction, or endocrine function."<sup>21</sup>

Nevertheless, because the types of safety-related research that can be conducted in humans are limited, animal studies are frequently cited in support of potential adverse effects. Results of these studies may be of questionable value due to the many physiological differences between animals and humans. Furthermore, many animals, including rodents and monkeys, metabolize isoflavones very differently than humans.<sup>62</sup> For a review of some of the key issues, see reference.<sup>63</sup> In 2006, the National Toxicology Program (NTP) Center for the Evaluation of Risks to Human Reproduction

evaluated the safety of soy formula. Although their initial conclusions supported the safety of soy formula use, no final report was issued.<sup>64, 65</sup> In 2009, the NTP again took up this issue. The conclusion of the 14-member panel of independent scientists was that there was "minimal concern" (the five levels of concern are negligible concern, minimal concern, some concern, concern and serious concern) about the safety of soy formula. Two panel members dissented from this consensus opinion, one in favor of "negligible concern" and the other in support of "some concern."

### **Effects of Soy Protein on Cholesterol Levels in Children**

As with adults,<sup>66</sup> clinical research in children shows that soy protein directly lowers serum cholesterol levels and improves levels of other lipids.<sup>67-71</sup> In the most recent study, when soy protein (average intake 0.5 g/kg body weight) was incorporated into the diets of children and adolescents (mean age 8.8 years; range 4-18 years) with familial and polygenic hypercholesterolemia, low density lipoprotein cholesterol decreased by 6.4 percent beyond the 11 percent decrease that occurred in response to the adoption of standard low-saturated fat diet during the three-month run-in period.<sup>71</sup> Thus, soy protein used in combination with other dietary therapies may reduce cholesterol levels to target goals.<sup>72</sup>

Soy protein may also serve as an adjunct to therapy in children taking medication for lowering cholesterol, there by reducing the required dose which may help to minimize or eliminate side effects.

### **Soy Protein Quality**

Soyfoods provide high-quality protein and are generally low in saturated fat.<sup>73</sup> Soy protein can meet protein needs of growing children. In 2000, the U.S. Department of Agriculture removed limits on the amount of soy protein that can be used in the National School Lunch Program.<sup>74</sup>

Providing healthful sources of protein without excessive saturated fat content is important for children. Higher protein diets are associated with greater satiety and weight loss.<sup>75</sup> Also, recent evidence in young boys shows that consumption of protein above



the recommended dietary allowance enhances the favorable impact of physical activity on bone mineral density.<sup>76</sup>

However, many protein-rich foods in children's diets are high in saturated fat. Therefore, substituting soyfoods for more traditional sources of protein generally improves overall diet quality. Even substituting soy protein for part of the beef or pork protein in a recipe can lead to a decrease in the fat, saturated fat and calorie content for the total entree, as long as portion size stays the same.<sup>77</sup> Similarly, combining cheese, eggs, or meat with tofu leads to improved nutritional quality of entrees.<sup>79</sup>

In general, soyfoods help children meet the Dietary Guidelines.<sup>77</sup> Short-term studies show that soyfoods support the normal growth and development of children<sup>80</sup> and improve growth when substituted for legumes in the diets of malnourished preschoolers.<sup>81</sup> Thus, soyfoods can play an important part in a healthy and varied diet.

### Acceptance of Soyfoods in Children's Diets

Research shows that soyfoods are generally well-accepted by children.<sup>79, 83, 84</sup> For example, among preschool children aged three to six years who attended a Head Start program, children consumed soy-enhanced lunches as readily as those made with more traditional ingredients, as evidenced by the amounts eaten.<sup>83</sup>

Negative beliefs about soy's palatability persist among some populations, however. When non-vegetarian subjects were told that a product contained soy, they were more likely to rate it as "grainy, chalky, dry, and unappealing" even when the product did not actually contain any soy ingredients.<sup>85</sup> Foods containing soy are also generally thought by U.S. consumers to be more "healthy tasting."<sup>85</sup> Ratings reflect the amount of soy consumed by a given individual.

### Soy Protein and Allergies

Essentially all food proteins have the potential to cause allergic reactions in some individuals. Although soy protein is one of the eight food proteins responsible for approximately 90 percent of all allergic reactions, these eight foods are not equally allergenic. The number of adults allergic to soy is quite small.<sup>86</sup> The relative number of children allergic to soy protein is almost certainly higher than the number of adults because children are much more sensitive to

*Soyfoods are generally well-accepted by children according to studies.*

dietary proteins.<sup>87</sup> However, most children are thought to outgrow their soy allergies early on in life,<sup>87</sup> although the pace at which this occurs is a matter of some recent discussion.<sup>88</sup> One recent study reported that more than 80 percent of infants outgrew their soy allergy by two years of age.<sup>89</sup>

### Isoflavones in Children's Diets

Preliminary data suggest that children actually absorb isoflavones to a greater extent than adults.<sup>90</sup> Although soyfoods have been consumed by Asian children for centuries without any apparent adverse effects, there is much interest in understanding the biological effects of isoflavones in children. In this regard, an Australian study published in 2008 examined the effects of isoflavones on high-density lipoprotein (HDL) levels<sup>91</sup> in boys. HDL levels decrease in boys as they enter puberty whereas no such decrease occurs in girls, a difference that may be due to the higher estrogen levels in females. Thus, it was hypothesized that isoflavone exposure would raise HDL levels in boys. However, no such changes occurred; thus, at least for this one possible measure of estrogenicity, no effects were noted. In agreement, research shows that soy protein has no effect on circulating testosterone<sup>92</sup> or estrogen levels in men.<sup>93-101</sup> Also as noted previously, no hormonal effects were noted in infants fed soy formula in the study most capable of detecting such changes.

Finally, there are speculative although very intriguing epidemiologic<sup>102-105</sup> and animal<sup>106, 107</sup> data suggesting that soy intake when young reduces breast cancer risk later in life. This evidence is consistent with mounting data that early life events greatly impact breast cancer risk.<sup>108</sup> The first twenty years of life appear to be particularly important.<sup>109</sup>



*Data suggests that soy intake during adolescence reduces breast cancer risk later in life.*

Research from the University of Alabama has shown that when rats are given the primary isoflavone in soybeans for just a few weeks early in life and then put on a typical laboratory diet, they develop 50 percent fewer tumors than rats not given this isoflavone.<sup>106</sup> These studies show that isoflavone exposure causes mammary cells to be changed in a way that makes them permanently less likely to be transformed into cancer cells later in life.<sup>110</sup> The protective effects of early pregnancy appear to work through a similar mechanism.<sup>108</sup>

*The potential public health benefit of modest soy consumption during childhood and adolescence cannot be overstated.*

The epidemiologic data suggest that quite modest amounts (perhaps just one serving) of soy during the early years are likely sufficient to reduce breast cancer risk.<sup>102-105</sup> The period of exposure to soy that is theoretically most protective against breast cancer is unclear. Although most studies have focused on the teenage years,<sup>102, 103, 105</sup> the results from a small study by Korde et al.<sup>104</sup> suggest soy consumption during childhood may be even more protective.

### Summary and Conclusions

Establishing good eating habits early in life is important. Childhood dietary intake may impact adult chronic disease risk and influence eating habits in adulthood. Soyfoods provide important options for improving the diets of young people, and research shows that these foods are acceptable to children.

Therefore, soyfoods can be viewed as healthy additions to the diets of children and adolescents. Other than relatively rare soy protein allergy, there is no clinical evidence that soyfoods exert any adverse effects. To the contrary, there is evidence suggesting that exposure to soy during childhood and/or adolescence reduces breast cancer risk later in life.

### References

1. Mikkila V, Rasanen L, Raitakari OT, Pietinen P, Viikari J. Consistent dietary patterns identified from childhood to adulthood: the cardiovascular risk in Young Finns Study. *Br J Nutr* 2005;93:923-31.
2. Mennella JA, Griffin CE, Beauchamp GK. Flavor programming during infancy. *Pediatrics* 2004;113:840-5.
3. Mennella JA, Beauchamp GK. Flavor experiences during formula feeding are related to preferences during childhood. *Early Hum Dev* 2002;68:71-82.
4. Birch LL. Development of food acceptance patterns in the first years of life. *Proc Nutr Soc* 1998;57:617-24.
5. Anonymous. Guidelines for school health programs to promote lifelong healthy eating. Centers for Disease Control and Prevention. *MMWR Recomm Rep* 1996;45:1-41.
6. Adair LS, Prentice AM. A critical evaluation of the fetal origins hypothesis and its implications for developing countries. *J Nutr* 2004;134:191-3.
7. McCormack VA, dos Santos Silva I, De Stavola BL, Mohsen R, Leon DA, Lithell HO. Fetal growth and subsequent risk of breast cancer: results from long term follow up of Swedish cohort. *BMJ* 2003;326:248.
8. Robinson R. The fetal origins of adult disease. *BMJ* 2001;322:375-6.
9. Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH. Long-term morbidity and mortality of overweight adolescents. A follow-up of the Harvard Growth Study of 1922 to 1935. *N Engl J Med* 1992;327:1350-5.
10. Russo J, Lareef H, Tahin Q, Russo IH. Pathways of carcinogenesis and prevention in the human breast. *Eur J Cancer* 2002;38 Suppl 6:S31-2.
11. Ogden CL, Carroll MD, Curtin LR, McDowell MA, Tabak CJ, Flegal KM. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA* 2006;295:1549-55.
12. Sorof JM. Prevalence and consequence of systolic hypertension in children. *Am J Hypertens* 2002;15:575-605.
13. Lipton RB. Incidence of diabetes in children and youth--tracking a moving target. *JAMA* 2007;297:2760-2.
14. Berenson GS, Srinivasan SR, Bao W, Newman WP, 3rd, Tracy RE, Wattigney WA. Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults. The Bogalusa Heart Study. *N Engl J Med* 1998;338:1650-6.
15. Breastfeeding and the use of human milk. American Academy of Pediatrics. Work Group on Breastfeeding. *Pediatrics* 1997;100:1035-9.
16. Ahluwalia IB, Morrow B, Hsia J, Grummer-Strawn LM. Who is breast-feeding? Recent trends from the pregnancy risk assessment and monitoring system. *J Pediatr* 2003;142:486-91.
17. Merritt RJ, Jenks BH. Safety of soy-based infant formulas containing isoflavones: the clinical evidence. *J Nutr* 2004;134:1220S-4S.
18. Osborn DA, Sinn J. Soy formula for prevention of allergy and food intolerance in infants. *Cochrane Database Syst Rev* 2006:CD003741.

19. Bhatia J, Greer F. Use of soy protein-based formulas in infant feeding. *Pediatrics* 2008;121:1062-8.
20. Kemp AS, Hill DJ, Allen KJ, et al. Guidelines for the use of infant formulas to treat cows milk protein allergy: an Australian consensus panel opinion. *Med J Aust* 2008;188:109-12.
21. Van Wyk JJ, Arnold MB, Wynn J, Pepper F. The effects of a soybean product on thyroid function in humans. *Pediatrics* 1959;24:752-60.
22. Shepard TH, Gordon EP, Kirschvink JF, McLean CM. Soybean goiter. *New Engl J Med* 1960;262:1099-103.
23. Pinchera A, MacGillivray H, Crawford JD, Freeman AG. Thyroid refractiveness in an athyreotic cretin fed soybean formula. *N Engl J Med* 1965;273:83-7.
24. Conrad SC, Chiu H, Silverman BL. Soy formula complicates management of congenital hypothyroidism. *Arch Dis Child* 2004;89:37-40.
25. Messina M, Redmond G. Effects of soy protein and soybean isoflavones on thyroid function in healthy adults and hypothyroid patients: a review of the relevant literature. *Thyroid* 2006;16:249-58.
26. Klein KO. Isoflavones, soy-based infant formulas, and relevance to endocrine function. *Nutr Rev* 1998;56:193-204.
27. Lasekan JB, Ostrom KM, Jacobs JR, et al. Growth of newborn, term infants fed soy formulas for 1 year. *Clin Pediatr (Phila)* 1999;38:563-71.
28. American Academy of Pediatrics. Committee on Nutrition. Soy protein-based formulas: recommendations for use in infant feeding. *Pediatrics* 1998;101:148-53.
29. American Academy of Pediatrics. Committee on Nutrition. Hypoallergenic infant formulas. *Pediatrics* 2000;106:346-9.
30. Barrett JR. Soy and children's health: a formula for trouble. *Environ Health Perspect* 2002;110:A294-6.
31. Barrett JR. The science of soy: what do we really know? *Environ Health Perspect* 2006;114:A352-8.
32. Setchell KD, Zimmer-Nechemias L, Cai J, Heubi JE. Exposure of infants to phytoestrogens from soy-based infant formula. *Lancet* 1997;350:23-7.
33. Barrett JR. The science of soy: what do we really know? *Environ Health Perspect* 2006 Jun;114:A352-8.
34. Setchell KD, Zimmer-Nechemias L, Cai J, Heubi JE. Exposure of infants to phytoestrogens from soy-based infant formula. *Lancet*. 1997;350:23-7.
35. Ho JY, Chen MJ, Sheu WH, Yi YC, Tsai AC, Guu HF, Ho ES. Differential effects of oral conjugated equine estrogen and transdermal estrogen on atherosclerotic vascular disease risk markers and endothelial function in healthy postmenopausal women. *Hum Reprod*. 2006 Oct;21:2715-20.
36. Lakoski SG, Brosnihan B, Herrington DM. Hormone therapy, C-reactive protein, and progression of atherosclerosis: data from the Estrogen Replacement on Progression of Coronary Artery Atherosclerosis (ERA) trial. *Am Heart J*. 2005 Nov;150:907-11.
37. Helgason S, Damber JE, Damber MG, von Schoultz B, Selstam G, Sodergard R. A comparative longitudinal study on sex hormone binding globulin capacity during estrogen replacement therapy. *Acta Obstet Gynecol Scand*. 1982;61:97-100.
38. Serin IS, Ozcelik B, Basbug M, Aygen E, Kula M, Erez R. Long-term effects of continuous oral and transdermal estrogen replacement therapy on sex hormone binding globulin and free testosterone levels. *Eur J Obstet Gynecol Reprod Biol*. 2001 Dec 1;99:222-5.
39. Reid IR, Eastell R, Fogelman I, Adachi JD, Rosen A, Netelenbos C, Watts NB, Seeman E, Ciaccia AV, Draper MW. A comparison of the effects of raloxifene and conjugated equine estrogen on bone and lipids in healthy postmenopausal women. *Arch Intern Med*. 2004 Apr 26;164:871-9.
40. Shulman LP. Effects of progestins in different hormone replacement therapy formulations on estrogen-induced lipid changes in postmenopausal women. *Am J Cardiol*. 2002 Jun 20;89:47E-54E; discussion E-5E.
41. Marqusee E, Braverman LE, Lawrence JE, Carroll JS, Seely EW. The effect of droloxifene and estrogen on thyroid function in postmenopausal women. *J Clin Endocrinol Metab*. 2000 Nov;85:4407-10.
42. Abech DD, Moratelli HB, Leite SC, Oliveira MC. Effects of estrogen replacement therapy on pituitary size, prolactin and thyroid-stimulating hormone concentrations in menopausal women. *Gynecol Endocrinol*. 2005 Oct;21:223-6.
43. Davies GC, Huster WJ, Shen W, Mitlak B, Plouffe L, Jr., Shah A, Cohen FJ. Endometrial response to raloxifene compared with placebo, cyclical hormone replacement therapy, and unopposed estrogen in postmenopausal women. *Menopause*. 1999 Fall;6:188-95.
44. Meuwissen JH, van Langen H. Monitoring endometrial thickness during estrogen replacement therapy with vaginosonography. *Radiology*. 1992 Apr;183:284.
45. Kaari C, Haidar MA, Junior JM, Nunes MG, Quadros LG, Kemp C, Stavale JN, Baracat EC. Randomized clinical trial comparing conjugated equine estrogens and isoflavones in postmenopausal women: a pilot study. *Maturitas*. 2006 Jan 10;53:49-58.
46. Yildiz MF, Kumru S, Godekmerdan A, Kutlu S. Effects of raloxifene, hormone therapy, and soy isoflavone on serum high-sensitive C-reactive protein in postmenopausal women. *Int J Gynaecol Obstet*. 2005 Aug;90:128-33.
47. D'Anna R, Baviera G, Corrado F, Cancellieri F, Crisafulli A, Squadrito F. The effect of the phytoestrogen genistein and hormone replacement therapy on homocysteine and C-reactive protein level in postmenopausal women. *Acta Obstet Gynecol Scand*. 2005 May;84:474-7.
48. Garrido A, De la Maza MP, Hirsch S, Valladares L. Soy isoflavones affect platelet thromboxane A2 receptor density but not plasma lipids in menopausal women. *Maturitas*. 2006 Jun 20;54:270-6.
49. Hall WL, Vafeiadou K, Hallund J, Bugel S, Reimann M, Koebnick C, Zunft HJ, Ferrari M, Branca F, et al. Soy-isoflavone-enriched foods and markers of lipid and glucose metabolism in postmenopausal women: interactions with genotype and equol production. *Am J Clin Nutr*. 2006 Mar;83:592-600.
50. Katz DL, Evans MA, Njike VY, Hoxley ML, Nawaz H, Comerford BP, Sarrel PM. Raloxifene, soy phytoestrogens and endothelial function in postmenopausal women. *Climacteric*. 2007 Dec;10:500-7.
51. Cheng G, Wilczek B, Warner M, Gustafsson JA, Landgren BM. Isoflavone treatment for acute menopausal symptoms. *Menopause*. 2007 May-Jun;14:468-73.
52. Bruce B, Messina M, Spiller GA. Isoflavone supplements do not affect thyroid function in iodine-replete postmenopausal women. *J Med Food*. 2003 Winter;6:309-16.
53. Marini H, Minutoli L, Polito F, Bitto A, Altavilla D, Atteritano M, Gaudio A, Mazzaferro S, Frisina A, et al. Effects of the phytoestrogen genistein on bone metabolism in osteopenic postmenopausal women: a randomized trial. *Ann Intern Med*. 2007 Jun 19;146:839-47.
54. Sarmmartino A, Di Carlo C, Mandato VD, Bifulco G, Di Stefano M, Nappi C. Effects of genistein on the endometrium: ultrasonographic evaluation. *Gynecol Endocrinol*. 2003 Feb;17:45-9.
55. Khaodhiar L, Ricciotti HA, Li L, Pan W, Schickel M, Zhou J, Blackburn GL. Daidzein-rich isoflavone aglycones are potentially effective in reducing hot flashes in menopausal women. *Menopause*. 2008 Jan-Feb;15:125-32.
56. Setchell KD. Assessing risks and benefits of genistein and soy. *Environ Health Perspect*. 2006 Jun;114:A332-3.
57. Munro IC, Harwood M, Hlywka JJ, Stephen AM, Doull J, Flamm WC, Adlercreutz H. Soy isoflavones: a safety review. *Nutr Rev*. 2003 Jan;61:1-33.
58. Gilchrist JM, Moore MB, Andres A, Estroff JA, Badger TM. Ultrasonographic Patterns of Reproductive Organs in Infants Fed Soy Formula: Comparisons to Infants Fed Breast Milk and Milk Formula. *J Pediatr*. 2009 Oct 19.
59. Strom BL, Schinnar R, Ziegler EE, Barnhart KT, Sammel MD, Macones GA, Stallings VA, Drulis JM, Nelson SE, Hanson SA. Exposure to soy-based formula in infancy and endocrinological and reproductive outcomes in young adulthood. *JAMA*. 2001 Aug 15;286:807-14.
60. Boucher BA, Cotterchio M, Kreiger N, Thompson LU. Soy Formula and Breast Cancer Risk. *Epidemiology*. 2008 Jan;19:165-6.
61. Chen A, Rogan WJ. Isoflavones in soy infant formula: a review of evidence for endocrine and other activity in infants. *Annu Rev Nutr*. 2004;24:33-54.
62. Gu L, House SE, Prior RL, Fang N, Ronis MJ, Clarkson TB, Wilson ME, Badger TM. Metabolic Phenotype of Isoflavones Differ among Female Rats, Pigs, Monkeys, and Women. *J Nutr*. 2006 May;136:1215-21.
63. Badger TM, Gilchrist JM, Pivik RT, Andres A, Shankar K, Chen JR, Ronis MJ. The health implications of soy infant formula. *Am J Clin Nutr*. 2009 May;89:1668S-72S.
64. Rozman KK, Bhatia J, Calafat AM, Chambers C, Culty M, Etzel RA, Flaws JA, Hansen DK, Hoyer PB, et al. NTP-CERHR expert panel report on the reproductive and developmental toxicity of genistein. *Birth Defects Res B Dev Reprod Toxicol*. 2006 Dec;77:485-638.
65. Rozman KK, Bhatia J, Calafat AM, Chambers C, Culty M, Etzel RA, Flaws JA, Hansen DK, Hoyer PB, et al. NTP-CERHR expert panel report on the reproductive and developmental toxicity of soy formula. *Birth Defects Res B Dev Reprod Toxicol*. 2006 Aug;77:280-397.
66. Zhan S, Ho SC. Meta-analysis of the effects of soy protein containing isoflavones on the lipid profile. *Am J Clin Nutr*. 2005 Feb;81:397-408.
67. Laurin D, Jacques H, Moorjani S, Steinke FH, Gagne C, Brun D, Lupien PJ. Effects of a soy-protein beverage on plasma lipoproteins in children with familial hypercholesterolemia. *Am J Clin Nutr*. 1991 Jul;54:98-103.
68. Widhalm K, Brazda G, Schneider B, Kohl S. Effect of soy protein diet versus standard low fat, low cholesterol diet on lipid and lipoprotein levels in children with familial or polygenic hypercholesterolemia. *J Pediatr*. 1993 Jul;123:30-4.
69. Gaddi A, Descovich GC, Nosedà G, Fragiaco C, Nicolini A, Montanari G, Vanetti G, Sirtori M, Gatti E, Sirtori CR. Hypercholesterolaemia treated by soybean protein diet. *Arch Dis Child*. 1987 Mar;62:274-8.
70. Blumenschein S, Torres E, Kushmaul E, Crawford J, Fixler D. Effect of oat bran/soy protein in hypercholesterolemic children. *Ann N Y Acad Sci*. 1991;623:413-5.
71. Weghuber D, Widhalm K. Effect of 3-month treatment of children and adolescents with familial and polygenic hypercholesterolaemia with a soya-substituted diet. *Br J Nutr*. 2008 Feb;99:281-6.
72. Jenkins DJ, Kendall CW, Marchie A, Faulkner DA, Wong JM, de Souza R, Emam A, Parker TL, Vidgen E, et al. Effects of a dietary portfolio of cholesterol-lowering foods vs lovastatin on serum lipids and C-reactive protein. *JAMA*. 2003 Jul 23;290:502-10.
73. Rand WM, Pellett PL, Young VR. Meta-analysis of nitrogen balance studies for estimating protein requirements in healthy adults. *Am J Clin Nutr*. 2003 Jan;77:109-27.
74. U.S. Department of Agriculture. Modification of the Vegetable Protein Products Requirements for the National School Lunch Program, School Breakfast Program, Summer Food Service Program and Child and Adult Care Food Program. *Fed Regist*. 2000;7 CFR Parts 210, 215, 220, 225 and 226:12429-42.

75. Astrup A. The satiating power of protein--a key to obesity prevention? *Am J Clin Nutr.* 2005 Jul;82:1-2.
76. Chevalley T, Bonjour JP, Ferrari S, Rizzoli R. High-Protein Intake Enhances the Positive Impact of Physical Activity on BMC in Prepubertal Boys. *J Bone Miner Res.* 2008 Jan;23:131-42.
77. Thomas JM, Lutz SF. Soy protein lowers fat and saturated fat in school lunch beef and pork entrees. *J Am Diet Assoc.* 2001 Apr;101:461-3.
78. McMIndes MK. Applications of isolated soy protein in low-fat meal products. *Food Technology.* 1991;45:61-4
79. Ashraf HR, Schoeppel C, Nelson JA. Use of tofu in preschool meals. *J Am Diet Assoc.* 1990 Aug;90:1114-6.
80. Egana JI, Fuentes A, Steinke FH, Uauy R. Protein quality comparison of a new isolated soy protein and milk in Chilean preschool children. *Nutr Res.* 1983;3:195-202.
81. Kay T, Ifeacho CL, Onowu G, Udenze V, Nnamani G, Anazonwu-Bello JN, Ifenu F, Modebe V, Emembolu M. Use of soya bean to improve the protein content of the diet in West Africa and thus prevent kwashiorkor. *J Trop Pediatr Environ Child Health.* 1975 Feb;21:45-8.
82. Mathew A, Raut DS. Effect of soymilk on the growth of malnourished children admitted to hospital wards. *Ind J Nutr Dietet.* 1981;18:260-7.
83. Endres J, Barter S, Theodora P, Welch P. Soy-enhanced lunch acceptance by preschoolers. *J Am Diet Assoc.* 2003 Mar;103:346-51.
84. Reilly JK, Lanou AJ, Barnard ND, Seidl K, Green AA. Acceptability of soymilk as a calcium-rich beverage in elementary school children. *J Am Diet Assoc.* 2006 Apr;106:590-3.
85. Wansink B, Westgren R. Profiling taste-motivated segments. *Appetite.* 2003 Dec;41:323-7.
86. Vierk KA, Koehler KM, Fein SB, Street DA. Prevalence of self-reported food allergy in American adults and use of food labels. *J Allergy Clin Immunol.* 2007 Jun;119:1504-10.
87. Cordle CT. Soy protein allergy: incidence and relative severity. *J Nutr.* 2004 May;134:12135-9S.
88. Skripak JM, Matsui EC, Mudd K, Wood RA. The natural history of IgE-mediated cow's milk allergy. *J Allergy Clin Immunol.* 2007 Nov;120:1172-7.
89. Aaronov D, Tasher D, Levine A, Somekh E, Serour F, Dalal I. Natural history of food allergy in infants and children in Israel. *Ann Allergy Asthma Immunol.* 2008 Dec;101:637-40.
90. Halm BM, Ashburn LA, Franke AA. Isoflavones from soya foods are more bioavailable in children than adults. *Br J Nutr.* 2007 Jul 11:1-8.
91. Dwyer T, Hynes KL, Fryer JL, Blizzard CL, Dalais FS. The lack of effect of isoflavones on high-density lipoprotein cholesterol concentrations in adolescent boys: a 6-week randomised trial. *Public Health Nutr.* 2008 Sep;11:955-62.
92. Hamilton-Reeves JM, Vazquez G, Duval SJ, Phipps WR, Kurzer MS, Messina MJ. Clinical studies show no effects of soy protein or isoflavones on reproductive hormones in men: results of a meta-analysis. *Fertil Steril.* 2009 Jun 11.
93. Tanaka M, Fujimoto K, Chihara Y, Torimoto K, Yoneda T, Tanaka N, Hirayama A, Miyanaga N, Akaza H, Hirao Y. Isoflavone supplements stimulated the production of serum equol and decreased the serum dihydrotestosterone levels in healthy male volunteers. *Prostate Cancer Prostatic Dis.* 2009 Jul 14.
94. Hamilton-Reeves JM, Rebello SA, Thomas W, Slaton JW, Kurzer MS. Isoflavone-Rich Soy Protein Isolate Suppresses Androgen Receptor Expression without Altering Estrogen Receptor- $\beta$  Expression or Serum Hormonal Profiles in Men at High Risk of Prostate Cancer. *J Nutr.* 2007 Jul;137:1769-75.
95. Goldin BR, Brauner E, Adlercreutz H, Ausman LM, Lichtenstein AH. Hormonal response to diets high in soy or animal protein without and with isoflavones in moderately hypercholesterolemic subjects. *Nutr Cancer.* 2005;51:1-6.
96. Dillingham BL, McVeigh BL, Lampe JW, Duncan AM. Soy protein isolates of varying isoflavone content exert minor effects on serum reproductive hormones in healthy young men. *J Nutr.* 2005 Mar;135:584-91.
97. Gardner-Thorpe D, O'Hagen C, Young I, Lewis SJ. Dietary supplements of soya flour lower serum testosterone concentrations and improve markers of oxidative stress in men. *Eur J Clin Nutr.* 2003 Jan;57:100-6.
98. Mitchell JH, Cawood E, Kinniburgh D, Provan A, Collins AR, Irvine DS. Effect of a phytoestrogen food supplement on reproductive health in normal males. *Clin Sci (Lond).* 2001 Jun;100:613-8.
99. Nagata C, Takatsuka N, Shimizu H, Hayashi H, Akamatsu T, Murase K. Effect of soymilk consumption on serum estrogen and androgen concentrations in Japanese men. *Cancer Epidemiol Biomarkers Prev.* 2001 Mar;10:179-84.
100. Higashi K, Abata S, Iwamoto N, Ogura M, Yamashita T, Ishikawa O, Ohszu F, Nakamura H. Effects of soy protein on levels of remnant-like particles cholesterol and vitamin E in healthy men. *J Nutr Sci Vitaminol (Tokyo).* 2001 Aug;47:283-8.
101. Habito RC, Montalto J, Leslie E, Ball MJ. Effects of replacing meat with soyabean in the diet on sex hormone concentrations in healthy adult males. *Br J Nutr.* 2000 Oct;84:557-63.
102. Shu XO, Jin F, Dai Q, Wen W, Potter JD, Kushi LH, Ruan Z, Gao YT, Zheng W. Soyfood Intake during Adolescence and Subsequent Risk of Breast Cancer among Chinese Women. *Cancer Epidemiol Biomarkers Prev.* 2001 May;10:483-8.
103. Wu AH, Yu MC, Tseng CC, Stanczyk FZ, Pike MC. Dietary patterns and breast cancer risk in Asian American women. *Am J Clin Nutr.* 2009 Feb 11.
104. Korde LA, Wu AH, Fears T, Nomura AM, West DW, Kolonel L, Pike MC, However R, Ziegler RG. Childhood soy intake and breast cancer risk in Asian American women. *Cancer Epidemiol Biomarkers and Prev.* 2009;18:1-9.
105. Lee SA, Shu XO, Li H, Yang G, Cai H, Wen W, Ji BT, Gao J, Gao YT, Zheng W. Adolescent and adult soy food intake and breast cancer risk: results from the Shanghai Women's Health Study. *Am J Clin Nutr.* 2009 Jun;89:1920-6.
106. Lamartiniere CA, Zhao YX, Fritz WA. Genistein: mammary cancer chemoprevention, in vivo mechanisms of action, potential for toxicity and bioavailability in rats. *J Women's Cancer.* 2000;2:11-9.
107. Peng JH, Zhang F, Zhang HX, Fan HY. Prepubertal octylphenol exposure up-regulate BRCA1 expression, down-regulate ER $\alpha$  expression and reduce rat mammary tumorigenesis. *Cancer Epidemiol.* 2009 Jul;33:51-5.
108. Russo J, Mailo D, Hu YF, Balogh G, Sheriff F, Russo IH. Breast differentiation and its implication in cancer prevention. *Clin Cancer Res.* 2005 Jan 15;11:931s-6s.
109. Shimizu H, Ross RK, Bernstein L, Yatani R, Henderson BE, Mack TM. Cancers of the prostate and breast among Japanese and white immigrants in Los Angeles County. *Br J Cancer.* 1991;63:963-6.
110. Messina M, Hilakivi-Clarke L. Early intake appears to be the key to the proposed protective effects of soy intake against breast cancer *Nutr Cancer.* 2009;61:792-8.



The United Soybean Board (USB) is a farmer-led organization comprised of 68 farmer-directors. Working with independent academic researchers affiliated with the National Institutes of Health (NIH) and academic institutions, USB has invested millions of dollars into health and nutrition research related to soy. Soybean farmers take pride in producing one of the healthiest food crops in the world. To access healthy soy recipes and more nutrition information, please visit [SoyConnection.com](http://SoyConnection.com).